## Method for Transforming Continuous PM2.5 Monitoring Data for Comparison with the FRM



## Michael Rizzo<sup>a</sup>, Peter Scheff<sup>b</sup>, and William Kaldy<sup>c</sup>

<sup>a</sup>USEPA Region 5, Air Monitoring

<sup>b</sup>University of Illinois — Chicago School of Public

Health, Environmental and Occupational

Health Sciences

<sup>c</sup>Hamilton County, OH Department of Environmental Services

## Data

- Looked at a variety sites from across country
  - All TEOMs
  - Variations in methodology
    - Season
    - Correction factors
- Sites with more than 1 year's worth of data
  - Collocated PM2.5 FRM and continuous method
  - Needed at least 63 samples across entire year to create model and R<sup>2</sup> at least 0.77
    - Based on USEPA DQOs
  - "Surplus" data used to validate model


## Data

- Valid days had 18 out of 24 hours
  - All variables had to meet requirement
- Model development
  - Hourly PM2.5 averaged over 24 hours
  - 24 hour FRM measurment
  - 24 hour average temperature
    - Collected from FRM monitor


## Validation

- National Weather Service temperatures
  - 2000 through 2001
- Closest met station temperatures
  - **2002**
- Examined hourly transformations averaged over 24 hours
  - Compared to FRM


#### WINTER PARK, FL




#### WINSTON-SALEM, NC




#### INDIANAPOLIS, IN



#### GRAND RAPIDS, MI



#### NEW YORK CITY, NY



## Methodology

#### "Knot" Method

**Equation 1** (if the temperature is less than the value of the knot)

```
FRM = \beta_0 + \beta_1*(avetemp-knot) + \beta_3*cont + \beta_4*cont(avetemp-knot) where:
```

**FRM** is the Federal Reference Method measurement

 $\beta_0$  is the intercept

 $\beta_1$  is the coefficient for the temperature term for temperatures less than the knot avetemp is the daily average temperature

*knot* is the temperature at which the linear relationship between the FRM and continuous measurement changes

 $\beta_3$  is the coefficient for the continuous measurement

**cont** is the continuous TEOM measurement

 $\beta_4$  is the coefficient of the interaction between the TEOM and temperature measurements for temperature less than the knot

# 4

## Methodology

#### "Knot" Method

**Equation 2** (if the temperature is greater than the value of the knot)

FRM =  $\beta_0$  +  $\beta_2$ \*(avetemp-knot) +  $\beta_3$ \*cont +  $\beta_5$ \*cont(avetemp-knot) where:

**FRM** is the Federal Reference Method measurement

 $\beta_0$  is the intercept

 $\beta_2$  is the coefficient for the temperature term for temperatures less than the knot

**avetemp** is the daily average temperature

*knot* is the temperature at which the linear relationship between the FRM and continuous measurement changes

 $\beta_3$  is the coefficient for the continuous measurement

**cont** is the continuous TEOM measurement

 $\beta_5$  is the coefficient of the interaction between the TEOM and temperature

measurements for temperature less than the knot

## Methodology

#### Linear Model

- FRM =  $\beta_0$  +  $\beta_1$ \*cont +  $\beta_2$ \*spring +  $\beta_3$ \*summer +  $\beta_4$ \*fall +  $\beta_5$ \*cont\*spring +  $\beta_6$ \*cont\*summer +  $\beta_7$ \*cont\*fall
- FRM = Federal Reference Method
- Cont = 24 hour avg continuous measurement
- Spring, summer, fall = seasonal variables
- Cont\*spring, cont\*summer, cont\*fall = interaction terms

# Results "Knot" Model

- Continuous parameter (β<sub>3</sub>) usually close to 1
- Temperature>"Knot" interaction term  $(\beta_5)$  usually not statistically significant
- Temperature<"Knot" interaction term</li>
   (β<sub>4</sub>) usually statistically significant
- Knot
  - Median: 15.4° C




## Results Seasonal Linear Model

- R<sup>2</sup> comparable to "Knot" model
- Surrogate for temperature in "Knot" model
- Disadvantage
  - Changes in season from fitting data change relationship between FRM and continuous measurement




# Examples of Model Fits Using Data Models Constructed From


#### NEW YORK CITY, NY




#### INDIANAPOLIS, IN




### CINCINNATI, OH




#### GRAND RAPIDS, MI




## ST PETERSBURG, FL




#### CHARLOTTE, NC



#### FORT WORTH, TX




## KEOSAUQUA, IA






## Examples of Validation Using "Extra" Data


#### NEW YORK CITY, NY




#### INDIANAPOLIS, IN




#### CINCINNATI, OH




#### GRAND RAPIDS, MI




## ST PETERSBURG, FL




#### CHARLOTTE, NC




#### FORT WORTH, TX



## KEOSAUQUA, IA



#### 2001 Fine Particulate Speciation Concentrations



### Conclusions

- "Knot" and seasonal linear method satisfy DQO requirements
- Some sort of seasonal adjustment necessary in areas with nitrate problem
- "Knot" and seasonally adjusted linear models comparable
  - Linear model surrogate for "Knot" model
- Uncertainty about seasonally adjusted linear model under changing conditions

## Conclusions

- Need for more data
  - Other monitoring technologies
  - Validation purposes
- Need for consistent national operating procedures across methodologies is **ESSENTIAL**
- Statistical transformation is temporary solution
  - Need technological solution implemented consistently across country
    - No change in data by use of various transformations across States
    - Better ensures data consistency and quality across States